

LIGO and Virgo make first detection of gravitational waves produced by colliding neutron stars

LIGO gii Virgo idumanistsduumya noo?gutsiisaduumya Abuduuxbiisii o?bigimskAAsts i_dubitsiibya nAbuutsbii bugutsitsbiixiya gagadusiix

Discovery marks first cosmic event observed in both gravitational waves and light.

oonoyiiyaatsimya ita?dawaa maduum?t idabobo?gwissxiniip muustsk Abuduuxbiisii
o?bigimskAAsts gii aanagimaa.

Watch the Press Conference live at: []

gidaxsatsiibuwa ayak?sessduuya Adudus: []

For the first time, scientists have directly detected gravitational waves — ripples in space-time — in addition to light from the spectacular collision of two neutron stars. This marks the first time that a cosmic event has been viewed in both gravitational waves and light.

aguduumaaniistsii, mugagyabiguwAx agA~gamu?gutsiisaduumya Abuduuxbiisii
o?biginskAAsts - gyanimI issxgwiip idAsiinimya - gyaniiaanagmaatsin uudo?batsuutsii
nadugIya bugutsitsbiixiya gagadusiix. anni ita?dawaa Aduum?stsii gyamu?k spuu?ts o?ganabii
itsatsimya miistsk Abuduuxbiisii o?bigimskAAsts gii aanagimaa

The discovery was made using the U.S.-based Laser Interferometer Gravitational-Wave Observatory (LIGO); the Europe-based Virgo detector; and some 70 ground- and space-based observatories.

gyamu?k oonoyiiyaatsimya itu?gwixxinimya amsgabuu?ts - itsii Laser Interferometer
Gravitational-Wave Observatory (LIGO); anni Europe-itsii Virgo itu?gutsiisaduubyuyix; gii
igiitsigibuu xa?gui- gii tsaanitsiibyuu-itsii idabobo?gwistsiists.

Neutron stars are the smallest, densest stars known to exist and are formed when massive stars explode in supernovas. As these neutron stars spiraled together, they emitted gravitational waves that were detectable for about 100 seconds; when they collided, a flash of light in the form of gamma rays was emitted and seen on Earth about two seconds after the gravitational waves. In the days and weeks following the smashup, other forms of light, or electromagnetic radiation — including X-ray, ultraviolet, optical, infrared, and radio waves — were detected.

aganyuup bugutsitsbiixiya gagadusiix iixstunado?bugiya gii igu?gadAxbigliya gii idAstuwasiiya
AsAxgaasimiists miisk stunaduumma?gu gagadusiix, gidoda~kuu?biiya bugutsitsbiixiya
gagadusiix, idabsgibiya muustsk Abuduuxbiisii o?bigimskAAsts itu?gutsiisaduumya gii**bii**bu
issxiinidaxin; Abuutsbiiyasix, ni_gI stunatsi~gamanatsi nidAni_gaduumya spisuuumskuutsp

aanatsiists AsAxgaasimya idabsgibya gii inimya anuum xa?gui naduixisnidaxin Apstsiixisamuu miixsk Abuduuxbiisii o?bigimskAAsts. madagAixtsiigui udAbuutsbiiya, myanitstsiiya aanatsiists noo?gutsiisaduumya.

The observations have given astronomers an unprecedented opportunity to probe a collision of two neutron stars. For example, observations made by the U.S. Gemini Observatory, the European Very Large Telescope, and the Hubble Space Telescope reveal signatures of recently synthesized material, including gold and platinum, solving a decades-long mystery of where about half of all elements heavier than iron are produced.

amuustsk essxgatsimya essbummi_datsiya idabesspsabiix ixxowanitsii ma?gitu?gwixxinisaa Abuutsbiisiists bugutsitsbiixiya gagadusiix. gii aganyuup idabobo?gwistsiimaanists anistutsimya miix U.S. Gemini Observatory, miix European Very Large Telescope, gii NASA's Hubble Space Telescope Astamatsugii nistsisgI Anagu abAsdutsiibya nigudAgimya, anu?k agAsxiniip tsiya ano~gii nigudAgimiists igutsistssuguuyi miistsk miixgim i_tu?dabAsdutsibya.

The LIGO-Virgo results are published today in the journal *Physical Review Letters*; additional papers from the LIGO and Virgo collaborations and the astronomical community have been either submitted or accepted for publication in various journals.

auu?k miix LIGO-Virgo agAxisdaniisdutsimya agitsiinAbya muu?k sinaixin *Physical Review Letters*; gyaniistsii sinaixin miix LIGO gii Virgo collaborations gii miix axiibiisabesspsabiix ago?gutsiaya gyI agotsimya agitsinsinaya myanistsiinapsxgixin

"It is tremendously exciting to experience a rare event that transforms our understanding of the workings of the universe," says France A. Córdova, director of the National Science Foundation (NSF), which funds LIGO. "This discovery realizes a long-standing goal many of us have had, that is, to simultaneously observe rare cosmic events using both traditional as well as gravitational-wave observatories. Only through NSF's four-decade investment in gravitational-wave observatories, coupled with telescopes that observe from radio to gamma-ray wavelengths, are we able to expand our opportunities to detect new cosmic phenomena and piece together a fresh narrative of the physics of stars in their death throes."

"igAdamidagyabii amu?k imagaabi otsisdabitsiip manistabodagiyan amuu spuu?ts, " awaanii France A. Cordova, Iiitsisgimaan miim National Science Foundation (NSF), uutsbuummuguwa miix LIGO. "amu?k oonoyiiyaatsimsin Adabisamuu nidabessginiibinnaan, na?gatsitu*essxgatsibinnaan spuu?ts o?ganabiists i_tu*dogutsiaya Abuduuxbiisii-o?bigimskAAsts idabobo?gwistsiists. aniixI NSF Adabisamuu nidessbuummuginnya mistsk Abuduuxbiisii-o?bigimskAAsts idabobo?gwistsiists gyamustk i_dAbissabyup i_tudessxgamatsimya i_dAn~ni_gyup gimiixsk spisuumskuutsp aanatsiists AsAxgaasimya isxgibiists, anu?gIya Adabssugabii agidoo?gutsiisaduumyat sim spuu?ts idIyidagiya gii agitu*dAtsinigliya gagadusiix axiniigisaa."

A stellar sign

gagadusii Anagu

The gravitational signal, named GW170817, was first detected on Aug. 17 at 8:41 a.m. Eastern Daylight Time; the detection was made by the two identical LIGO detectors, located in Hanford, Washington, and Livingston, Louisiana. The information provided by the third detector, Virgo, situated near Pisa, Italy, enabled an improvement in localizing the cosmic event. At the time, LIGO was nearing the end of its second observing run since being upgraded in a program called Advanced LIGO, while Virgo had begun its first run after recently completing an upgrade known as Advanced Virgo.

amuu Abuduuxbiisii Iyiinapsgagii, itsin_gaduup GW170817, iduum?togutsiisaduumya idepu*gyagyup igitsigigubudui, udAdudu?s 8:41 xisganoduni Eastern Daylight Time; amu?k oo?gutsiisaduumii idayiinapsgutsiiyawa miix nadugIya LIGO itu?gutsiisaduubyuyix, itstsiiya miim Hanford, Washington, gii Livingston, Louisiana. gyani Iiinapsgoogowa essbumumuyIyA miix nuxgessti itu?gutsiisaduubyuyix, Virgo, idutstsii Pisa, Italy, idAsbuummuguwa ma?gidAsuxxinibuwa tsima itu?gabii amu?k o?ganabii. animI, LIGO agayaxisdabiisuumya utu?nadugisstsii essxgamiyyasiists udAgyogadAdabuu?suwa utsiitsgagoo mu?k itsin_gaduup Advanced LIGO, gyamixii Virgo bitsiiyumadabuuya utsiitsgagoo itsin_gaduumya Advanced Virgo.

The NSF-funded LIGO observatories were conceived, constructed, and operated by Caltech and MIT. Virgo is funded by the Istituto Nazionale di Fisica Nucleare (INFN) in Italy and the Centre National de la Recherche Scientifique (CNRS) in France, and operated by the European Gravitational Observatory. Some 1,500 scientists in the LIGO Scientific Collaboration and the Virgo Collaboration work together to operate the detectors and to process and understand the gravitational-wave data they capture.

annii NSF-isssbummuuts LIGO idabobo?gwistsiists no?i_tsiiximstaaya, nAbuutsduum, gii odwitsigaduumya miix Caltech gii MIT. Virgo udessbummuuguwa miixsk Istituto Nazionale di Fisica Nucleare (INFN) miim Italy gii miixsk Centre National de la Recherche Scientifique (CNRS) miim France. aganyuup/aniixI 1500 mugagyabiguwAx annii LIGO Scientific Collaboration gii miixsk Virgo Collaboration Abuudabodagiya ma?gidodwitsigaduumya muuxsk itu?gutsiisaduubyuyix gii ma?gidotsisdabidatsuwa gii a~ku?tsigiy muustsk Abuduuxbiisii o?bigimskAAsts agoostsiimaan eganudaduumya.

Each observatory consists of two long tunnels arranged in an L shape, at the joint of which a laser beam is split in two. Light is sent down the length of each tunnel, then reflected back in the direction it came from by a suspended mirror. In the absence of gravitational waves, the laser light in each tunnel should return to the location where the beams were split at precisely the same time. If a gravitational wave passes through the observatory, it will alter each laser beam's arrival time, creating an almost imperceptible change in the observatory's output signal.

gyagiido idabobo?gwis anni nistsii nadugaya ixinuuyii i_dAstata?duup aniststsIya nibwisi annimI utsidessgaatsiya annii napstsanatsi idayagiidootsiya. aanagimaa idessgitsduu?bii miistsi gyagiido i_dAstata?duup, matsidessku?biiya itu?tsanidabstsiiya awayaagiisaa annii asaabyatsis. igamu?iisaaniyiiwa Abuduuxbiisiiya o?bigimskAAsts annii napstsanatsi agu?gitsduup gyagiido i_dAstata?duup agidessgitsduu?bii itu?tsanidabstsiiya miistsk idabanatsi awayaagiisaaya agu?gitsduup. igamanistsiisi muustsk Abuduuxbiisii o?bigimskAAsts agitu?tsitsgu amu?k idabobo?gwis, aku?gitsii gyagiido napstsanatsi idabanatsi utsidudu?buwl, agabistuutsiip stunatu*bugitsii annii idabobo?gwisin Iyiinapsgagii.

On Aug. 17, LIGO's real-time data analysis software caught a strong signal of gravitational waves from space in one of the two LIGO detectors. At nearly the same time, the Gamma-ray Burst Monitor on NASA's Fermi space telescope had detected a burst of gamma rays. LIGO-Virgo analysis software put the two signals together and saw it was highly unlikely to be a chance coincidence, and another automated LIGO analysis indicated that there was a coincident gravitational wave signal in the other LIGO detector. Rapid gravitational-wave detection by the LIGO-Virgo team, coupled with Fermi's gamma-ray detection, enabled the launch of follow-up by telescopes around the world.

ani_k idepu*gyagyup igitsigubudui, nidabidudu agooxtsiimaan idessxgatsimya egamuuxdagyup eganudagitsimya miistsk iigIyiigu Iyiinapsgagiists niistsii Abuduuxbiisii o?bigimskAAsts i_~tududuya spuu?ts giniixi niduxgIya LIGO itu?gutsiisaduubyuyix. iduutstsii, nixii spisuumskuutsp aanatsiists AsAxgaasimya o?gwiixiniip miim NASA's Fermi i_dAbissabyup itu?gutsiisaduumyatsix spisuumskuutsp aanatsiists AsAxgaasimya. LIGO-Virgo idessxgatsimya egamuuxdagyup nAbutu?duumya muustsk Iyiinapsgagiists gii itsiinimya madaku?danistsii gii matstsigii no?isabaduup LIGO idessxgatsimya awAstamatsiyya a?gitstsiip axigamaanistsii Abuduuxbiisii o?bigimskAAsts namatsiik LIGO itu?gutsiisaduubyuyix. egamuu Abuduuxbiisii- o?bigimskAAsts oo?gutsiisaduumiix anniixi LIGO-Virgo nessbumi_daya, giniixi Fermi's spisuumskuutsp aanatsiists AsAxgaasimya oo?gutsiisaduumya, agitu~tubii agitu?wisxinimya i_dAbissabyupiists annuum xa?gui.

The LIGO data indicated that two astrophysical objects located at the relatively close distance of about 130 million light-years from Earth had been spiraling in toward each other. It appeared that the objects were not as massive as binary black holes — objects that LIGO and Virgo have previously detected. Instead, the inspiraling objects were estimated to be in a range from around 1.1 to 1.6 times the mass of the sun, in the mass range of neutron stars. A neutron star is about 20 kilometers, or 12 miles, in diameter and is so dense that a teaspoon of neutron star material has a mass of about a billion tons.

muu?k LIGO agooxtsiimaan Astamatsugii nadugaya spuu?tsiists Iyiniibiists itstsiiya aganyuup igusatstsii tsaanitsibisstsiiya annuum xa?gui agAsamuu odaguuyatsiya. iganistsiinatsi muustsk Iyiniibiists madu?guma?xtsi miistsk nadugisstsii sigooxgiya - Iyiniibiists miix LIGO gii Virgo noo?gutsiisaduumya. misxawa amuuxk Iyiniibiiss odaguuya awaniiya inagutsitsguum?xiimi nah nadus niitsisxga?binaduugi , stunaduma?xtsii idanisdAsiiya miistsii bugutsitsbiixiya gagadusiix. bugutsitsbiixiya gagadusiix agnyuup anistsinam bugwi, nitsxgwip naatsigubudui gii stunat?dexbigiiya igamu?dudagyup niduxge buxim innu?suyis miistsk bugutsitsbiixiya gagadusiix igagissagu aganyuup noi miisdagiists.

While binary black holes produce "chirps" lasting a fraction of a second in the LIGO detector's sensitive band, the Aug. 17 chirp lasted approximately 100 seconds and was seen through the entire frequency range of LIGO — about the same range as common musical instruments. Scientists could identify the chirp source as objects that were much less massive than the black holes seen to date.

miistsk nadugisstsii sigooxgiya nido?gumiya 'biixiini_gi' idaabigamuui miim LIGO itu?gutsiisaduubyuyixiists idawatsimabiiya, anni? idepu*gyagyup igitsigubudui 'biixiini_gi' nitsii giibiiuibui issxiinidaxin gii no?ganAnimya muustsk essanistasi miim LIGO - isxgwipyatsix idanisdAsiiya i_deni_gyubiists. mugagyabiguwAx itsiinimya miistsk biixiini_giists Iyiniibiists madu?igagoyatsigiyag agtisdotsstuup miistsk sigooxga anu?k.

“It immediately appeared to us the source was likely to be neutron stars, the other coveted source we were hoping to see — and promising the world we would see,” says David Shoemaker, spokesperson for the LIGO Scientific Collaboration and senior research scientist in MIT’s Kavli Institute for Astrophysics and Space Research. “From informing detailed models of the inner workings of neutron stars and the emissions they produce, to more fundamental physics such as general relativity, this event is just so rich. It is a gift that will keep on giving.”

“tsiiganistsiitsiibinaan a?ganistabiiya bugutsitsbiixiya gagadusiix, amuxk dabesssatsdubinnaan - gii **nidawani_binnaan** **aku?ganAniip**,” awanii David Shoemaker, uudAdumaguwaa udAbuwaduumuuguwa miix LIGO Scientific Collaboration gii **uuma?gina agabo?gwixiniip** mugagyabiguwan miix MIT’s Kavil Institute muuxsk Astrophysics and Space Research. “no?guutsp itsiinxin ido?gwissxiniip muustsk bugutsitsbiixiya gagadudix gii **abatseegiidiuyiists abAstutsimya**, giimiistsk **manabii** essatsimmya tsiimaii_diitstsii naduusi gii gagaduusi gyamuu?k bisaatsinsiimaan, muu?k no?ganabi **igaguutsabii**. **axdaamooduu?giidasin**.”

“Our background analysis showed an event of this strength happens less than once in 80,000 years by random coincidence, so we recognized this right away as a very confident detection and a remarkably nearby source,” adds Laura Cadonati, professor of physics at Georgia Tech and deputy spokesperson for the LIGO Scientific Collaboration. “This detection has genuinely opened the doors to a new way of doing astrophysics. I expect it will be remembered as one of the most studied astrophysical events in history.”

“isskuu?ts **nidessxgatsibinnaan** **dAstamatsugiinnaan** amuu?k o?ganabii **ido?gabiya** Adabisamuu egamsuuganistsii, **nisudamsxiniipbinnaan** isgadaksin oo?gutsiisaduumyatsix gii iixisadabi iduutstsiw uutsu?**guunimanuwa**,” nu?kdawanii Laura Cadonati, espssxininatsdu?gii muustsk essatsimmya tsiimaii_duutstsii naduusi gii gagaduusi miim Georgia Tech gii ?batsinaa uudAdumaguwaa udAbuwaduumuuguwa miix LIGO Scientific Collaboration. “amu?k oo?gutsiisaduumyatsix **igayinniipyga** **gitsimiists** **maaniwatstiists** agitu?dodabodagiya mustsk **axiibiisabesspsatsimmya** tsiimaii_duutstsii naduusi gii gagaduusi. **nistsiiximst** **aku?ganessxiniip** essxinimatsdutsiip muustsk **axiibiisabesspsatsimmya** tsiimaii_duutstsii naduusi gii gagaduusi o?ganabiists agessxu*dAbuyiya.”

Theorists have predicted that when neutron stars collide, they should give off gravitational waves and gamma rays, along with powerful jets that emit light across the electromagnetic spectrum. The gamma-ray burst detected by Fermi, and soon thereafter confirmed by the European Space Agency’s gamma-ray observatory INTEGRAL, is what’s called a short gamma-ray burst; the new observations confirm that at least some short gamma-ray bursts are generated by the merging of neutron stars — something that was only theorized before.

matsidabiix **Iyistamattsagiya** **iigayissxinimya** miistsk bugutsitsbiixiya gagadusiix agibuutxbiyyasa, agitstsiip Abuduuxbiisii o/bigimskAAsts gii spisuumskuutsp aanatsiists AsAxgaasimya, gyaniistsii ibigapssi skunadabi asiita?da **abatseesiiduumya myaanagimaatsiists**. mu?k spisuumskuutsp aanatsiists AsAxgaasimya oo?gutsiisaduumyatsix miix Fermi nidAn_gadaya magaispisuumskuutsp aanatsiists AsAxgaasimya; gyamuu **manessxgamaatsimya** agidugamuudanyuup gyamuustsii magaispisuumskuutsp aanatsiists AsAxgaasimya **idabistuutsiibya** mustsk agiibuutstu?biya bugutsitbiixiya gagadusiix - bitsiistuu Agaku?**dAbuyiya**.

“For decades we’ve suspected short gamma-ray bursts were powered by neutron star mergers,” says Fermi Project Scientist Julie McEnery of NASA’s Goddard Space Flight Center. “Now, with the incredible data from LIGO and Virgo for this event, we have the answer. The gravitational waves tell us that the merging objects had masses consistent with neutron stars, and the flash of gamma rays tells us that the objects are unlikely to be black holes, since a collision of black holes is not expected to give off light.”

“Adabisamu o?gutsisaaduup magaispisuumskuutsp aanatsiists AsAxgaasimya madumii babusiists niistsk bugutsitsbiixiya gagadusiix iibuut?biiya,” awanii Fermi Project Scientist Julie McEnery i_tu*dudu NASA’s Goddard Space Flight Center. “anu?k, ibisadabii agooxtsimaaniists idaisdudaya miix LIGO gii Virgo mi_k oo?ganabii, agessxiniip. muxsk Abuduuxbiisii o?bigimskAAsts awaniigiyia iibuut?biya Iyiniibiists igoma?guyiists niiduyaanistsiis mistsk bugutsitsbiixiya gagadusii , gii idaabigamuu napstsanatsi miistsk spisuumskuutsp aanatsiists AsAxgaasimya i_tu*tsxiniip miixsk Iyiniibiists madaganistabiya sigooxgiya, madesstums?kuutsiya aanagimaa.”

But while one mystery appears to be solved, new mysteries have emerged. The observed short gamma-ray burst was one of the closest to Earth seen so far, yet it was surprisingly weak for its distance. Scientists are beginning to propose models for why this might be, McEnery says, adding that new insights are likely to arise for years to come.

anistsii agitsigAstamanistasii no?gudAtsiip, manigAstamanistasii. nistsii nessxgaduutsiipy mustsk magaispisuumskuutsp aanatsiists AsAxgaasimya igiduutstsii num xa?gui udAnibuwaduum, gyaganyuup iixgo?matsgunadapsii. mugagyabiguwAx anu?k essatsimya maganistsii, McEnery awanii, gii issu?tsiigi aganistsii igagawuu agitu?gwissxinimya.

A patch in the sky

buku*nebistsi amuu spuu?ts

Though the LIGO detectors first picked up the gravitational wave in the United States, Virgo, in Italy, played a key role in the story. Due to its orientation with respect to the source at the time of detection, Virgo recovered a small signal; combined with the signal sizes and timing in the LIGO detectors, this allowed scientists to precisely triangulate the position in the sky. After performing a thorough vetting to make sure the signals were not an artifact of instrumentation, scientists concluded that a gravitational wave came from a relatively small patch in the southern sky.

miix LIGO itu?gutsiisaduubyuyix inoku*uutsimya Abuduuxbiisii o?bigimskAAsts anuum amsgabuu?ts, Virgo, miim Italy, anixI maduumstsii amu?k atsinigiip. utsitstsiipuwa, udoo?gutsiisaduumyatsix mi_k uutsu?guunimanuwa, Virgo no?gunimya buuk*Iyiinapsgagi; Abuutsdudaduumya amuusts Iyiinapsgagiists essxgaatsimya gii maniistsiisamsii miistsk LIGO itu?gutsiisaduubyuyix, o?gutxsinimya muxsk mugagyabiguwAx nuxgIya Iyu?simiya tsima itsstsii spuu?ts. Apstsiixisamuu idagubaduumya ma?gitu?gwissxiniibuwa muustsk Iyiinapsgagiists idudabii miistsk edexistsigum, mugagyabiguwAx nitsi_daya muustsk Abuduuxbiisii o?bigimskAAsts itu?dudu stuunatu?bugi anim amsgabuu?ts spuu?ts.

“This event has the most precise sky localization of all detected gravitational waves so far,” says Jo van den Brand of Nikhef (the Dutch National Institute for Subatomic Physics) and VU University Amsterdam, who is

the spokesperson for the Virgo collaboration. “This record precision enabled astronomers to perform follow-up observations that led to a plethora of breathtaking results.”

“gyamu?k o?ganabii ikusoboyapssimaduumya tsima isspuu?ts amuustsk oo?gutsiisaduumya Abuduuxbiisii o?bigimskAAsts anu?k,” awanii Jo van den Brand i~tududu Nikhef (the Dutch National Institute of Subatomic Physics) gii VU University Amsterdam, aniistapsii uudAdumaguwaa udAbuwaduumuuguwa miix Virgo collaboration. “amu?k soboyapssimsin essbumuuya muxsk idabesspsabiix ma?gidessxgatsibuwa itsibisaatstsiya agAxisdaniisdutsiip.”

“This result is a great example of the effectiveness of teamwork, of the importance of coordinating, and of the value of scientific collaboration,” adds EGO Director Federico Ferrini. “We are delighted to have played our relevant part in this extraordinary scientific challenge: Without Virgo, it would have been very difficult to locate the source of the gravitational waves

“i~gamsawitstsii miix Virgo, igaxiiyigu a?gido?guniip uutsu?guunimanuwa mustsk Abuduuxbiisii o?bigimskAAsts,” nuu?gadawanii idAnaawasii miix European Gravitational Observatory Federico Ferrini.

Fermi was able to provide a localization that was later confirmed and greatly refined with the coordinates provided by the combined LIGO-Virgo detection. With these coordinates, a handful of observatories around the world were able, hours later, to start searching the region of the sky where the signal was thought to originate. A new point of light, resembling a new star, was first found by optical telescopes. Ultimately, about 70 observatories on the ground and in space observed the event at their representative wavelengths.

Fermi no?gu~tu*gutsiim tsima itu?duduyiists gidagidugamuudanyupa gii itsugabistutsimya miistsk isxgwibiists no?gutsiya nAgiibuutsduusiix LIGO-Virgo oo?gutsiisaduumya. udAsxinibuuwa miistsk isxgwibiists, itstsiya idabobo?gwistsiists anuum xa?gui, matsisamu, idumattsatsimya miim spuu?ts a?~kutsiximsstaya a?ku*dutstsii muu?k Iyiinapsgagii. anii mannaanatsii, anastsim gagadusiwa, i_tu*tsinimya i_dAbissabyutsiists. iidaniststsii, agnyuup i_gitsigbui idabobo?gwistsiists anuum xa?gui gii spuu?ts nessxgatsimya mi_k o?ganabii mixisk egmuuxdagyp oxstsimya.

“This detection opens the window of a long-awaited ‘multi-messenger’ astronomy,” says Caltech’s David H. Reitze, executive director of the LIGO Laboratory. “It’s the first time that we’ve observed a cataclysmic astrophysical event in both gravitational waves and electromagnetic waves — our cosmic messengers. Gravitational-wave astronomy offers new opportunities to understand the properties of neutron stars in ways that just can’t be achieved with electromagnetic astronomy alone.”

“amu?k oo?gutsiisaduumya igayinniip manii xistsiiguumsdan Adabiisamu?gitsiip ‘myanissxxsimiwa uududanigiyaa’ axiibiisabesspsabii,” awanii Caltech’s David H. Reitze, oxgasatsduum miix LIGO Laboratory. “manessxgatsibinaan amu?k uma?xsomiidababii axiibiisabesspsatsimmya tsimaii_duutstsii naduusi gii gagaduudiix o?ganabii niduysiists Abuduuxbiisii o?bigimskAAsts gii babusuuxbiisiists o?bigimskAAsts - amuxk spuu?ts uududanigiyaa Abuduuxbiisii o?bigimskAAsts axiibiisabesspsabii o?guguya manixxowanitsii agidotsistaibiitsiip amistapsiiya manisti_bii muxsk bugutsitsbiixiya gagadusiix igagiguu agitu?dabodagyup niitsk babusuuxbiisiists axiibiisabesspsabii niitstsiiwa.

A fireball and an afterglow

bagoyitsibugwi gii Apstsiixisamuu ixistui Apstsiixisamuu egaanagimaa

Each electromagnetic observatory will be releasing its own detailed observations of the astrophysical event. In the meantime, a general picture is emerging among all observatories involved that further confirms that the initial gravitational-wave signal indeed came from a pair of inspiraling neutron stars.

amuustsk **babusuxbiisiists** idabobo?gwis aku*Asesduuyatsimya utu?dessxgamaabitsim muu?k **axiibiisabesspsatsimmya** tsimaii_duutstsii naduusi giigagaduusi o?ganabii. aganyup, **agumadapsugessxinimya** Asaxstiya muustsk idabobo?gwistsiists **agAsuxxinimya** gyamu?k ixidui bitsisdui Abuduuxbiisii-o?bigimskAAsts Iyiinapsgagii i_~tududu misstsk nadugaya **nAsaguuya** bugutsitsbiixiya gagadusii.

Approximately 130 million years ago, the two neutron stars were in their final moments of orbiting each other, separated only by about 300 kilometers, or 200 miles, and gathering speed while closing the distance between them. As the stars spiraled faster and closer together, they stretched and distorted the surrounding space-time, giving off energy in the form of powerful gravitational waves, before smashing into each other.

aganyup gibibunuxga ixsigaasi agessduuyiists, nadugIya bugutsitsbiixiya gagadusiix oda~ku*biiya, manistibuu?bii aganyup **nadugibibasin** gii Agyiistunatsiigamuu?biiya utsidessgaatsiya. amuustsk gagadusiists odaku*biyasiya gii nAginAbuutstsiiya, Asepi_gaduumya gii **matsbumabiiyaduum** amuu udaguu?ts spuu?ts-agAdudu, iyiistapiksduum esstuumskuutspduumya Abuduuxbiisii o?bigimskAAsts, gyanimI itsibuutsbiisiya.

At the moment of collision, the bulk of the two neutron stars merged into one ultradense object, emitting a “fireball” of gamma rays. The initial gamma-ray measurements, combined with the gravitational-wave detection, also provide confirmation for Einstein’s general theory of relativity, which predicts that gravitational waves should travel at the speed of light.

gyanii udo?batsuutsiiya amuustsk udu?ma?gagabiists naduge bugutsitsbiixiya gagadusiix idanisdapsiya niduxge stunatsixbigi Iyiniibiista, nawabattsAxgumya amuuyI ‘bagoyitsibugwi’ anistsi spisuumskuutsp aanatsiists AsAxgaasimya. annii ixidui bitsisdui spisuumskuutsp aanatsiists AsAxgaasimya issxgwiibiists, nAbuutsduumya miistsk Abuduuxbiisii-o?bigimskAAsts oo?gutsiisaduumya, matsidagidugamuudandumya nam Einstein’s bisaatsinsiimaan, iigayissxinim miistsk Abuduuxbiisii o?bigimskAAsts a?gidegamu*biiya muustsk aanagimAsts.

Theorists have predicted that what follows the initial fireball is a “kilonova” — a phenomenon by which the material that is left over from the neutron star collision, which glows with light, is blown out of the immediate region and far out into space. The new light-based observations show that heavy elements, such as lead and gold, are created in these collisions and subsequently distributed throughout the universe.

matsidabiix Iyiist**mattsagiya** iiyissxinimya **matsigyabiwa** ammyI bagoyitsibugwi anniistapsii “**stunatsuma?guunAxgaasi gagadusii**” - omAdagisin aganyup amuustsk itsimatsiists namidutsitstxtsii annii bugutsitsbiixiya gagadusiix nAbuutsbiiya, egaanagimatsi, **nayistapu*biitsiisaduupya** amii spuu?ts. amustsk maaniya essxgamiisttsiists **Astamatsuugiya**

miistsk igesssugiists nigudAgimya, aganyuup odaigim, idabAstutsiibya Abuutsbiiyatsigiya gii idAganitsip*dubya spuu?ts.

In the weeks and months ahead, telescopes around the world will continue to observe the afterglow of the neutron star merger and gather further evidence about various stages of the merger, its interaction with its surroundings, and the processes that produce the heaviest elements in the universe.

madagAixtsiigui issu?tsiigi aganistsii, i_dAbissabyubiists anuum xa?gui axdamessatsimya niistsk Apstsiixisamuu egaanagimaa amuustsk bugutsitsbiixiya gagadusiix iibuutu?biiya gii umuu?dudaya iitu?gunimaaya amuustsk myanistsiiaxtsino?dubopya miistsk iibuutu?biiya, udessinogugubimaasts udaguu?ts, gii manistabAstutsipyia igessugiya nigudAgimiists miim spuu?ts.

“When we were first planning LIGO back in the late 1980s, we knew that we would ultimately need an international network of gravitational-wave observatories, including Europe, to help localize the gravitational-wave sources so that light-based telescopes can follow up and study the glow of events like this neutron star merger,” says Caltech’s Fred Raab, LIGO associate director for observatory operations. “Today we can say that our gravitational-wave network is working together brilliantly with the light-based observatories to usher in a new era in astronomy, and will improve with the planned addition of observatories in Japan and India.”

“nidomadabitsii_daan LIGO isskuu?ts 1980’s, nitsxinibinaan nidaxdotu*gwiik_binnaan uugyabi uudaabsduudaxin Abuduuxbiisii-o?bigimskAAsts idabobo?gwistsiists, annuyi Europe, agitsuutuu?guunisuwa Abuduuxbiisii-o?bigimskAAsts uutsu?guunimanuw aamiistsk aanatsi-itsii i_dAbissabyup agidessxgatsimya amuustsk egaanagimaa o?ganabiists aganyuup muuxsk bugutsitsbiixiya gagadusiix iibuutu?biiya,” awanii Caltech’s Fred Raab LIGO essbumuuyi ooxgaduumsin idabobo?gwis abodagiists. “anuu?k xistigui aku*gudanyuup muuxk Abuduuxbiisii-o?bigimskAAsts udaabsduudaxin stunadAsugabudugidagiya gyanii aanatsi-itsiiidabobo?gwistsiists axigayanimatsutsiip manAxiibiisabesspsabii, gii agutsitsxugabii itsii_daansinix miix idabobo?gwistsiists itstsiiya Japan gii India.”

LIGO is funded by the [NSF](#), and operated by [Caltech](#) and [MIT](#), which conceived of LIGO and led the Initial and Advanced LIGO projects. Financial support for the Advanced LIGO project was led by the NSF with Germany ([Max Planck Society](#)), the U.K. ([Science and Technology Facilities Council](#)) and Australia ([Australian Research Council](#)) making significant commitments and contributions to the project.

LIGO uutsbumuguwa miix [NSF](#), gii odwitsigaduumya miix [Caltech](#) gii [MIT](#), i_tsiiximstaaya LIGO gii agomadabanistsiiduumya miix Initial gii Advanced LIGO uudabuudaxix. uutsbumuguwa miix Advanced LIGO uudabuudaxix agomadabanistsiiduumya mmix NSF gimiixi Germany ([Max Planck Society](#)), miix U.K. ([Science and Technology Facilities Council](#)) gii Australia ([Australian Research Council](#)) ixsugaganisdutsiimya gii itsbuum_daya amuu uudabuudaxix.

More than 1,200 scientists and some 100 institutions from around the world participate in the effort through the [LIGO Scientific Collaboration](#), which includes the GEO Collaboration and the Australian collaboration OzGrav. Additional partners are listed at <http://ligo.org/partners.php>

aganyuup 1200 mugagyabiguwAx i_~tududuya gyanuum xa?gui agitsinsuubiya gyaxigagimaduup miix LIGO Scientific Collaboration, annuyi miix GEO Collaboration gii miix Australian essdudabodagiya OzGrav. matsuix nessbumi_dAx itsiinaaya
<http://ligo.org/partners.php>.

The Virgo collaboration consists of more than 280 physicists and engineers belonging to 20 different European research groups: six from Centre National de la Recherche Scientifique (CNRS) in France; eight from the Istituto Nazionale di Fisica Nucleare (INFN) in Italy; two in the Netherlands with Nikhef; the MTA Wigner RCP in Hungary; the POLGRAW group in Poland; Spain with the University of Valencia; and the European Gravitational Observatory, EGO, the laboratory hosting the Virgo detector near Pisa in Italy, funded by CNRS, INFN, and Nikhef.

miix Virgo collaboration anni nistsii 280 myanitsxiininatstu?xiniix gii nAybuutsduumya o?ganaduumya tsaniiduu?tstsiiya European agabo?gwisxinimiixa: noi miim Centre National de la Recherche Scientifique (CNRS) miim France; nanisui i_~tududuya Istituto Nazionale di Fisica Nucleare (INFN) miim Italy; nadugum miim Netherlands muuxsk Nikhef; miix MTA Wigner RCP miim Hungary; miix POLGRAW oo?ganuuya miim Poland; Spain miix University of Valencia; gii miix European Gravitational Observatory, EGO, miix igayiguu madaagabuudaxix aku?ganuuya miix Virgo itu?gutsiisaduubyuyix iduutstsiw Pisa miim Italy, utsbummuuguwa miix CNRS, INFN, gii Nikhef.

###

Written by Jennifer Chu, MIT News Office

RELATED LINKS

Paper: "GW170817: Observation of gravitational waves from a binary neutron star merger."
** Paper will be available to read online at 10:00 AM EDT on Oct. 16, 2017.

MEDIA CONTACTS

Kimberly Allen, MIT
allenkc@mit.edu; +1 617-253-2702

Emily Velasco, Caltech
evelasco@caltech.edu; +1 626-395-6487

Jason Maderer, Georgia Tech
maderer@gatech.edu; +1 404-385-2966

Severine Perus, Virgo-EGO
severine.perus@ego-gw.it; +39 050 752 325

Aya Collins, National Science Foundation
acollins@nsf.gov; +1 703-292-7737

* This press release is translated into the Siksika (Blackfoot) language by Sharon Yellowfly (Siksika Nation) who has a Bachelor's of Arts in Anthropology, a background in linguistics, & who is fluent in the Siksika language. This translation was made from a Blackfoot Dictionary Yellowfly has written. Words are spelled phonetically with the English alphabet, but here are notes on vowels & other symbols used to make sounds unique to the Siksika language:

Vowels & Other Sounds

SIKSIKA

a
i
u
e
o

ENGLISH

father
eat
book
let
go

x - six

A - acorn

I - ice

? - glottal stop

- - as in 'he' but held a little longer

* - *who*

~ - (not quite a full glottal stop) as in '*cotton*'

[bold letters] - inflection